

COVID-19 Antibody Treatment Discharge Instructions

Today you received: bamlanivimab 700 mg
 other: _____

⊗ After your infusion, you should go home and rest.

To help stop the spread of COVID-19, stay home and away from others until your healthcare provider tells you it is safe to resume normal activities.

⊗ Bruising and slight discomfort at the IV site is common and should go away in a few days.

You may use a cold compress for comfort today if needed. After that, warmth (like a heating pad) can help heal bruising at the site.

If you notice changes such as **pain, redness, drainage, numbness or tingling** or and other concerning symptoms near your IV site, contact your healthcare provider.

⊗ Tell your healthcare provider if you have side effects*, including:

- Upset stomach (nausea, vomiting, or diarrhea)
- Itching, swelling, rash, or hives
- Dizziness or low blood pressure
- Changes in your heartbeat
- Any new or worsening symptoms

You can also report side effects to FDA MedWatch at www.fda.gov/medwatch or by calling 1-800-FDA-1088

**If you have any of these emergency warning signs*,
get medical attention right away:**

- Pain or pressure in the chest
- Trouble breathing
- Wheezing (noisy breathing that may sound musical or like whistling)
- New confusion
- Inability to wake or stay awake
- Bluish lips or face
- Swollen lips, face, or throat

*This is not a complete list of possible symptoms. Please call your medical provider for any other symptoms that are severe or concerning to you.