

Patient Name:	`	
DOB:	,	

ADUCANUMAB-AVWA (ADUHELM™) INFUSION ORDERS

1)	☐ G30.1 Alzheimer's ☐ G30.8 Other Alzheimer's ☐ G31.84 Mild cogn	eimer's disea	secondary F	$02.8 \times \text{code} \rightarrow$ \square Other:		1 Dementia with	nout behavioral disturbance n behavioral disturbance
	Gender: □ M	□ F	Height:		□ CM	Weight:	□ KG □ LB
2	Beta-amyloid pathol	logy confirmed	he following red			en met (please	e provide documentation) Result:
3)	Cognitive assessmen						Result:
サ	☐ MRI obtained prior Pre-Infusion: ✓		auneim therapy (with			oate:	Result:
	☑ Calculate aducan ☑ Dilute required v	new o		s actual weig	ht and dose		o not round dose.
	-					oride and infuse	over at least 60 minutes
	-		nding 0.2- or 0.22-r	micron in-line	filter.		over at least 60 minutes
	-		nding 0.2- or 0.22-r Treatment Num	micron in-line Iber	filter. Weig	nt-based Dose	over at least 60 minutes
	-		nding 0.2- or 0.22-r	nicron in-lind Iber Ision 2	filter.		over at least 60 minutes
	-		nding 0.2- or 0.22-r Treatment Num Infusion 1 and Infu	micron in-lind liber Ision 2 Ision 4	e filter. Weig	nt-based Dose mg/kg	over at least 60 minutes
	-		nding 0.2- or 0.22-r Treatment Num Infusion 1 and Infu Infusion 3 and Infu	micron in-line ber usion 2 usion 4 usion 6	weigter. Weig 1 3	mg/kg	e over at least 60 minutes
	using a sterile, lo	w protein-bii	Treatment Num Infusion 1 and Infu Infusion 3 and Infu Infusion 5 and Infu Infusion 7 and be	nicron in-line ber usion 2 usion 4 usion 6 yond	Weig 1 3 6 10	mg/kg mg/kg mg/kg mg/kg mg/kg	over at least 60 minutes
	using a sterile, lo	w protein-bi	Treatment Num Infusion 1 and Infu Infusion 3 and Infu Infusion 5 and Infu Infusion 7 and be occurs, stop infus	nicron in-line liber lision 2 lision 4 lision 6 lyond lision and tre	Weige 1 3 6 10 at per order	mg/kg mg/kg mg/kg mg/kg mg/kg ers/protocol as	
	using a sterile, low ✓ If infusion-relat ✓ Schedule treatme	ed reaction	Treatment Num Infusion 1 and Infu Infusion 3 and Infu Infusion 5 and Infu Infusion 7 and be occurs, stop infus	nicron in-line liber lision 2 lision 4 lision 6 lyond lision and tre days apart). C	Weig 1 3 6 10 at per order	mg/kg mg/kg mg/kg mg/kg ers/protocol as	clinically indicated.
	using a sterile, low ☑ If infusion-relat ☑ Schedule treatme ☐ Order e Post-Infusion: ☑ Educate pati	ed reaction ents every 4 xpires on	Treatment Num Infusion 1 and Infu Infusion 3 and Infu Infusion 5 and Infu Infusion 7 and be occurs, stop infus weeks (at least 21 o	nicron in-line liber lision 2 lision 4 lision 6 lyond lision and tre ldays apart). Coder expires a	Weig 1 3 6 10 at per order valid to	mg/kg mg/kg mg/kg mg/kg mg/kg ers/protocol as for one year unle	clinically indicated.
	using a sterile, low ☑ If infusion-relat ☑ Schedule treatme ☐ Order e Post-Infusion: ☑ Educate pati	ed reaction ents every 4 expires on	Treatment Num Infusion 1 and Infu Infusion 3 and Infu Infusion 5 and Infu Infusion 7 and be occurs, stop infus weeks (at least 21 c	micron in-line liber lision 2 lision 4 lision 6 lyond lision and tre days apart). Coder expires and line, dizziness, pelow	Weig Weig 1 3 6 10 at per order valid to the mausea, vis	mg/kg mg/kg mg/kg mg/kg ers/protocol as for one year unle treatments ion changes, or i	c clinically indicated.

