

Discharge Instructions

Today you received. SPEVIGO® (spesolimab-sbzo) 900 mg

 **After your infusion you can resume normal activities.**





Some people feel a bit tired and prefer to rest after their infusions, and others feel energized. It is safe to resume normal activities if you feel up to it – just be sure to listen to your body and rest when needed!

 **Bruising and slight discomfort at the IV site is common and should go away in a few days.**

You may use a cold compress for comfort today if needed. After that, warmth (like a heating pad) can help heal bruising. Be careful not to freeze or burn your skin!

If you notice changes such as **pain, redness, drainage, numbness or tingling** or and other concerning symptoms near your IV site, contact your healthcare provider.

Tell your healthcare provider or get emergency medical help right away if you have signs of an allergic reaction*, including:

-  Swollen face, eyelids, lips, mouth, tongue, or throat
-  Hives (raised, itchy areas of skin)
-  Trouble breathing, or tightness in the chest or throat
-  Any new or worsening symptoms

*This is not a complete list of possible symptoms. Please call your medical provider for any other symptoms that are severe or concerning to you.

 **Follow up with your GPP specialist.**

If your generalized pustular psoriasis (GPP) flare symptoms don't get better, your specialist may decide that you should receive another SPEVIGO treatment in one week.