

lecanemab-irmb**LEQEMBI****100 mg/mL****Dose Preparation**
**NATIONAL
INFUSION CENTER
ASSOCIATION**
Order Details**Patient Weight:**
*(enter patient weight in kg
and click on Tab key)*
kg**Dose @ 10 mg/kg****mg****Dose Volume:****mL****Diluent Volume:****mL**

0.9% Sodium Chloride

Total Volume:
 (Dose Volume + Diluent
Volume)
mL**Dose Range****500 mg / 5 mL****vials needed****200 mg / 2 mL****vials needed**

500 mg

1

0

501 - 700 mg

1

1

701 - 900 mg

1

2

901 - 1000 mg

2

0

1001 - 1200 mg

2

1

1201 - 1400 mg

2

2

1401 - 1500 mg

3

0

Dose Preparation Instructions

- 1 Calculate the number of LEQEMBI (lecanemab-irmb) vials needed for the dose (see above). Each vial contains a concentration of 100 mg / mL.
- 2 Remove LEQEMBI (lecanemab-irmb) vial(s) from refrigerated storage and allow to warm to room temperature. Do not expose to direct heat.
- 3 Visually inspect the LEQEMBI (lecanemab-irmb) diluted solution for particles or discoloration prior to administration. Do not use if it is discolored, or opaque or foreign particles are seen.
- 4 Remove the flip-off cap from the vial. **Disinfect and allow to air dry. Using aseptic technique**, insert the sterile syringe needle into the vial through the center of the rubber stopper.
- 5 Slowly withdraw _____ mL of LEQEMBI (lecanemab-irmb) from the vial. Each vial is for single-use only; discard any unused portion.
- 6 Disinfect injection port of infusion bag containing 250 mL of 0.9% Sodium Chloride Injection and allow to air dry. Using aseptic technique, slowly inject LEQEMBI (lecanemab-irmb) into infusion bag. Avoid creating turbulence or foaming.
- 7 Gently invert the infusion bag containing the LEQEMBI (lecanemab-irmb) diluted solution to mix completely. **Do not shake. Do not freeze.**
- 8 **After dilution, immediate use is recommended.** If not administered immediately, store LEQEMBI (lecanemab-irmb) refrigerated for up to 4 hours, or at room temperature for up to 4 hours.

Disclaimer: RxToolKit is providing this worksheet on an "AS IS" basis and the entire risk remains with the prescribing provider and clinicians administering care. Healthcare providers and clinicians prescribing or administering this medication should consult the most current medication package insert and/or prescribing information. RxToolKit disclaims any and all warranties or conditions. In no event will RxToolKit be liable for any damages, including without limitation any indirect, consequential, special, incidental or punitive damages arising out of, based on, or resulting from the use of RxToolKit materials. This worksheet is the copyright and intellectual property of RxToolKit.

lecanemab-irmb**LEQEMBI****Dose Administration****RxToolKit****NATIONAL
INFUSION CENTER
ASSOCIATION****Order Details**

| | |
|--|-----------|
| Dose: | mg |
| Dose Volume: | mL |
| Diluent Volume: 0.9% Sodium Chloride | mL |
| Total Volume: (Dose Volume + Diluent Volume) | mL |

Order Details

| | |
|----------------------------|--|
| Administration Time | 1 hour |
| Filter | 0.2 micron in-line filter |
| Flush | Yes <i>Flush infusion line to insure all drug is administered.</i> |
| Infusion Rate | mL/hr |

Administer by Intravenous Infusion

Attach the infusion set (containing a terminal low-protein binding 0.2 micron in-line filter) to the IV bag. Prime the infusion set.

Infuse the entire volume of the LEQEMBI (lecanemab-irmb) diluted solution intravenously over approximately one hour.

Flush infusion line to ensure all LEQEMBI (lecanemab-irmb) is administered.

Monitor for signs and symptoms of an infusion reaction.

The infusion rate may be reduced, or the infusion may be discontinued, and appropriate therapy administered as clinically indicated.

Consider pre-medication at subsequent dosing with antihistamines, non-steroidal anti-inflammatory drugs, or corticosteroids, if signs or symptoms of an infusion-related reaction are observed.